

This question paper contains 8 printed pages]

Code No. : 23(I)

Roll No.

0(CCEM)9

STATISTICS

Paper : I

Time Allowed : 3 hours]

[Maximum Marks : 300

Note : (i) Answers must be written in English.

(ii) Number of marks carried by each question are indicated at the end of the question.

(iii) Part/Parts of the same question must be answered together and should not be interposed between answers to other questions.

(iv) The answer to each question or part thereof should begin on a fresh page.

(v) Your answers should be precise and coherent.

(vi) Attempt five questions in all, choosing at most two questions from each Section. Question No. 1 is compulsory.

P. T. O.

SECTION - I

(Probability)

1. (a) Write down the three basic axioms of probability and show that if the events A and B are such that $A \subset B$ then :
- (i) $P(A) \leq P(B)$, and
- (ii) $P(B - A) = P(B) - P(A)$ 15
- (b) What are the various methods of estimation ? Explain any *two* of them with examples. 15
- (c) State and prove Neyman-Pearson lemma. 15
- (d) Describe the analysis of two-way classified data and give the ANOVA table. 15
2. (a) Define conditional probability and show that it satisfies axioms of probability. 20
- (b) Let A and B be the respective events that two contracts I and II are completed by certain deadlines and suppose that P (at least one contract is completed by the deadline) = 0.9 and P (both contracts are completed by their deadlines) = 0.5. Calculate the probability P (exactly one contract is completed by its deadline). 20
- (c) Let the events A_1, A_2, \dots, A_n defined on a sample space Ω be such that :

(i) $A_i \cap A_j = \phi, i \neq j,$

(ii) $\bigcup_{i=1}^n A_i = \Omega,$ and

(iii) $P(A_i) > 0$ for all i . Then for any event B on Ω , show that :

$$P(B) = \sum_{i=1}^n P(B/A_i) \cdot P(A_i) \quad 20$$

3. (a) Define random variable (r. v.) and its distribution function (d. f.). A continuous r. v. X has probability density function (pdf) :

$$f(x) = \frac{2x}{9}, 0 < x < 3$$

$$= 0, \text{ otherwise.}$$

Find :

(i) d. f. of X

(ii) $P(X \leq 2)$

(iii) $P(-1 < x < 1.5)$

(iv) $P(X = 2)$

20

(3)

P. T. O.

23(I)

- (b) Calculate the characteristic function (ch. f.) of a r. v. X with pdf :

$$f(x) = \lambda e^{-\lambda x}, x \geq 0$$

Hence obtain its mean and variance. 20

- (c) A bowl contains 8 chips. Three of the chips are red and remaining 5 are blue. Two chips are drawn successively at random and without replacement. What is the probability that the first draw results in a red chip and the second draw results in a blue chip? 20

4. (a) Given that the joint pdf of r. vs. X and Y is

$$f(x, y) = \frac{x(1+3y^2)}{4}, 0 < x < 2$$

$$= 0, \quad \text{otherwise}$$

Find :

- (i) The marginal pdfs of X and Y ,
(ii) The conditional pdf of X given $Y = y$,
(iii) $P\left(\frac{1}{4} < X < \frac{1}{2} \mid Y = \frac{1}{3}\right)$ 20

- (b) Explain the weak and strong law of large numbers. 20

- (c) Let $u(X)$ be a non-negative function of r. v. X .
If $E(u(X))$ exists then for every constant C ,
show that :

$$P(u(X) \geq C) \leq \frac{E(u(X))}{C}$$

Hence deduce Chebyshev's inequality. 20

SECTION - II

(Statistical Inference)

5. (a) Define consistent estimator of a parameter. Show that in random sampling from a normal population, the sample mean is a consistent estimator of the population mean. 30
- (b) Show that $x(x-1)/n(n-1)$ is an unbiased estimator of θ^2 for the $B(n, \theta)$ distribution. 30
6. (a) How is Cramer-Rao inequality useful in obtaining minimum variance unbiased estimator (MVUE) ?
Given the pdf :

$$f(x; \theta) = \frac{e^{-\theta} \cdot \theta^x}{x!}, x = 0, 1, 2, \dots; \theta > 0$$

Show that the C-R lower bound is θ/n . 30

23(I)

- (b) Describe sequential probability ratio test (SPRT), operating characteristic (OC) and Average sample number (ASN) functions. Derive an expression for ASN. Also obtain ASN for testing $H_0 : \mu = 10$ against $H_1 : \mu = 20$, where observations follow $N(\mu, 10)$ distribution. (Given $\alpha = 0.04$, $\beta = 0.10$). 30

7. (a) What is meant by confidence interval? How are confidence intervals obtained for large samples? Show that for the distribution:

$$f(x; \theta) = \theta \cdot e^{-\theta x}, \quad x \geq 0$$

the confidence limits for θ in large samples with confidence coefficient $\alpha = 0.05$ are given by:

$$\left(1 \pm \frac{1.96}{\sqrt{n}} \right) / \bar{x} \quad 30$$

- (b) Define sufficient and efficient estimator. Given that X_1 and X_2 are two independent observations from a Poisson distribution with mean θ , show that $X_1 + X_2$ is sufficient for θ and $X_1 + 2X_2$ is not sufficient for θ . 30

8. (a) State and prove generalized Neyman Pearson lemma for randomized test: 30

- (b) Let θ be a real parameter and let X have pdf:

$$f(x, \theta) = c(\theta) \cdot \exp \{Q(\theta)T(x)\} h(x),$$

(6)

where $Q(\theta)$ is strictly monotone. Prove that for testing $H_0 : \theta \leq \theta_0$ against $H_1 : \theta > \theta_0$, there exists a UMPU test ϕ such that, if $Q(\theta)$ is increasing then :

$$\begin{aligned}\phi(x) &= 1 & \text{if } T(x) > c \\ &= \gamma & \text{if } T(x) = c \\ &= 0 & \text{if } T(x) < c,\end{aligned}$$

where c and γ are obtained by :

$$E_{\theta_0} \phi(x) = \alpha. \quad 30$$

SECTION - III

(Linear Inference & Multivariate Analysis)

9. (a) Define estimable function and show that a parametric function $l' \beta$ is estimable if and only if l can be expressed as a linear combination of rows of X . 30
- (b) Define Hotelling's T^2 -statistic. Show that :

$$\lambda^{2/n} = \frac{1}{1 + \frac{T^2}{n-1}}$$

where symbols have their usual meaning. 30

10. (a) For the general linear model $\underline{y} = X\underline{\beta} + \underline{e}$, where X is of order $n \times p$ ($p < n$) matrix, show that there exist no unbiased estimator of $\underline{\beta}$ if $\text{rank}(X) < p$. 30

23(I)

- (b) Define multiple correlation coefficient R between x_1 and (x_2, x_3, \dots, x_p) and show that $R^2 \geq R_0^2$, where R_0 is multiple correlation coefficient between x_1 and any linear combination of x_2, x_3, \dots, x_p . 30
11. (a) Suppose $\underline{X} \sim N_p(\underline{\mu}, \Sigma)$ and let \underline{X} be partitioned as $\underline{X} = (\underline{X}'_1, \underline{X}'_2)$, with $\text{Cov}(\underline{X}_1, \underline{X}_2) = \Sigma_{12}$. Prove that the necessary and sufficient condition for \underline{X}_1 and \underline{X}_2 to be independent is that $\Sigma_{12} = 0$. 30
- (b) Discuss discriminant analysis with reference to the case of two p -variate normal populations. How will you test the goodness of an assigned discriminant function? 30
12. (a) Discuss the method of fitting an orthogonal polynomials. What are the advantages of using orthogonal polynomials for fitting curvilinear relations? 30
- (b) Define Hotelling's T^2 statistic. Give its applications in testing (i) the significance of mean vector of a multivariate normal distribution; (ii) the equality of mean vectors of two multivariate normal populations. 30